

Neotropical *Nilothauma* Kieffer, 1921 (Diptera, Chironomidae): Key, eleven new species, re-descriptions, new combination and new records

Luiz Carlos Pinho¹, Trond Andersen²

1 Laboratory of Systematics of Diptera, Department of Ecology and Zoology, Federal University of Santa Catarina, Campus Trindade, CEP 88040-900, Florianópolis, Brazil **2** Department of Natural History, University Museum of Bergen, University of Bergen, P.O. Box 7800, NO-5020, Bergen, Norway

Corresponding author: Luiz Carlos Pinho (luiz.pinho@ufsc.br)

Academic editor: F.L. da Silva | Received 11 November 2020 | Accepted 1 February 2021 | Published 22 April 2021

<http://zoobank.org/BEFB015F-4A35-4987-AD10-DFEB7D4F91DD>

Citation: Pinho LC, Andersen T (2021) Neotropical *Nilothauma* Kieffer, 1921 (Diptera, Chironomidae): Key, eleven new species, re-descriptions, new combination and new records. ZooKeys 1033: 81–125. <https://doi.org/10.3897/zookeys.1033.60686>

Abstract

Nine new species of *Nilothauma* Kieffer, *N. hamadae* sp. nov., *N. jupau* sp. nov., *N. karitiana* sp. nov., *N. leccii* sp. nov., *N. marianoii* sp. nov., *N. mateusi* sp. nov., *N. txukuyana* sp. nov., *N. werekena* sp. nov. and *N. yekwana* sp. nov. are described and figured, based on adult males collected in Brazil and *N. maya* sp. nov. on an adult male from Mexico; *N. terena* sp. nov. is described as male, pupa and larva based on a reared specimen from Brazil. *Nilothauma aleta* Roback, 1960 and *N. duena* Roback, 1960 are re-described and recorded from Brazil. *Nilothauma longissimum* Mendes & Andersen, 2009 is transferred to *Beardius* Reiss & Sublette, 1985 and the diagnosis of *Nilothauma* is emended. New records of thirteen Neotropical *Nilothauma* species are given and a key to the males of all known species of *Nilothauma* is provided.

Keywords

Brazil, key, Mexico, Neotropical Region, new combination, new records, new species, taxonomy

Introduction

The genus *Nilothauma* was erected by Kieffer (1921), based on the Afrotropical *N. picipenne* Kieffer, 1921. The Neotropical genera *Neelamia* Sponis, 1987 and *Paranilothauma* Sponis, 1987 were both placed as synonyms of *Nilothauma* by Mendes and Andersen (2009). Most males of *Nilothauma* can be recognised by having an antenna with 13 flagellomeres, low antennal ratio (except in *Nilothauma longissimum* Mendes & Andersen, 2009), fore-tibia with long spur on conical, apical scale, high venarum ratio and squama bare (Mendes and Andersen 2009). In addition, many males have median or lateral, sometimes strongly setose lobes or projections on tergite IX. The larvae of *Nilothauma* inhabit littoral and sublittoral soft sediments of lakes, streams and rivers (Epler et al. 2013).

To date, the genus comprises 52 described species that occur in all zoogeographical regions, except Antarctica (Qi et al. 2014, 2016; Niitsuma 2016; Dantas and Hamada 2017). A total of 22 species are known from the Neotropical and four from the Nearctic Regions (Andersen et al. 2016; Dantas and Hamada 2017).

After examining material collected in several localities in the Neotropical Region, mostly from the Brazilian Amazon, 24 species of *Nilothauma* were identified. Eleven of them proved to be new to science and are described below as adult males and for *N. terena* sp. nov. also as larva and pupa. The other thirteen species have their distribution range extended in the Neotropics. *Nilothauma aleta* Roback, 1960 and *N. duena* Roback, 1960, originally described from the Peruvian Amazon are re-described, based on material from Brazil and *N. longissimum* Mendes & Andersen, 2009 is transferred to *Beardius* Reiss & Sublette, 1985. A key to the males of all known species of *Nilothauma* is provided.

Material and methods

Alcohol-preserved specimens were dissected and slide-mounted in Euparal. Morphological terminology and abbreviations follow Sæther (1980). Measurements are taken according to Epler (1988) and given as ranges, followed by the mean when more than three specimens were measured, followed by the number of specimens measured in parenthesis.

Abbreviations used in the text as follows:

- CEPA** Centro de Estudos e Pesquisas Ambientais (Centre for Environmental Studies and Research);
- EB** Estação Biológica (Biological Station);
- INPA** Instituto Nacional de Pesquisas da Amazônia (Manaus, Brazil);
- MZSP** Museu de Zoologia da Universidade de São Paulo (São Paulo, Brazil);
- PE** Parque Estadual (State Park);
- RPPN** Reserva Particular do Patrimônio Natural (Private Natural Heritage Reserve);

- UFSC** Entomological Collection of the Federal University of Santa Catarina (Florianópolis, Brazil);
ZMBN University Museum of Bergen (Bergen, Norway);
ZSM Zoologische Staatssammlung München (Munich, Germany).

Type material is deposited at INPA, MZSP, UFSC, ZMBN and ZSM, as stated in each description. Vouchers are deposited at UFSC, ZMBN and ZSM.

Taxonomy

Beardius Reiss & Sublette, 1985

Beardius longissimus (Mendes & Andersen, 2009), **comb. nov.**

Nilothauma longissimum Mendes & Andersen, 2009: 26

Material examined. Type material, as in Mendes and Andersen (2009).

Additional material. 8 males, slide-mounted: BRAZIL, Bahia, Camacan, RPPN Serra Bonita, Trilha Bapeba, 15°20'35"S, 39°33'34"W, 4.xi.2009, light trap, A.R. Calor et al. leg. 3 males, slide-mounted, as previous, except: 15°23'32"S, 39°33'53"W, 2.xi.2009. 1 male, slide-mounted, as previous, except: 03.ii.2009. 1 male, slide-mounted, as previous, except: córrego 2, 15°23'10"S, 39°34'03"W, 819 m a.s.l., 01.viii.2008, light trap, A.R. Calor, L.S. Lecci, L.C. Pinho & R.A. Moretto leg. 1 male, slide-mounted: BRAZIL, São Paulo, PE Serra do Mar, Ubatuba, Picinguaba, Camburi stream, 09.ix.2006, light trap, M.R. Spies & A.E. Sieglöch leg.

Remarks. Pinho et al. (2013) found that the “presence of apical thin setae on inferior volsella” (character 74, state 1), i.e. a group of two, rarely three, slender simple setae at the very tip of the inferior volsella, in addition to subapical, stouter setae, is the only synapomorphy of *Beardius* Reiss & Sublette, 1985 in the adult stage. In fact, the character is shared by all species of *Beardius* and is not found elsewhere, except in *Nilothauma longissimum* Mendes & Andersen, 2009, a species that was considered to be sister to all remaining *Nilothauma* by Mendes and Andersen (2009) in a phylogenetic analysis with *Paratendipes* Kieffer and *Pseudochironomus* Malloch as outgroups. The comparatively-high antennal ratio in *N. longissimum* (AR > 1.00) is the only exception in *Nilothauma*, which generally have very low antennal ratios (AR < 0.40). Further, the venarum ratio (VR) seems to be low in *N. longissimum* compared to other *Nilothauma* species; other characters in *N. longissimum* are consistent with the current diagnosis of *Beardius*. We therefore propose the new combination and emend the diagnosis of *Nilothauma* accordingly.

Distribution. The species was originally described from São Paulo State, south-eastern Brazil by Mendes and Andersen (2009); the range is now extended to Bahia State in north-eastern Brazil.

Nilothauma Kieffer, 1921

Emended diagnosis. After transferring *Nilothauma longissimum* Mendes & Andersen, 2009 to *Beardius* Reiss & Sublette, 1985 (see above), adult males of *Nilothauma* become more easily separated from other genera. The diagnosis given by Mendes and Andersen (2009) has to be emended as follows: “antennal ratio generally low (AR < 0.40), one species (*N. longissimum* sp. nov.) with AR > 1.00.” should read: “antennal ratio generally low (AR < 0.40), occasionally as high as 0.82 (*N. soka* Andersen, Bello González & Hagenlund, 2016).

The discovery of the pupae of *N. terena* sp. nov. leads to the diagnosis of the pupa given by Mendes and Andersen (2009) having to be emended as follows: “Frontal setae short, not on tubercles.” should read “Frontal setae short, occasionally long and taeniate, not on tubercles.” Further, “Sternites I–VII bare; sternite VIII with central, longitudinal field of shagreen.” should read “Sternites I–VII usually bare, sternite I occasionally with extensive shagreen; sternite VIII with central, longitudinal field of shagreen.”

Based on the larva of *N. terena* sp. nov., the diagnosis of the *Nilothauma* larvae in Epler et al. (2013) should be emended as follows: “Mandible. All teeth pale;” should read: “Mandible. All teeth pale, occasionally inner teeth with somewhat darker pigmentation;” and “Mentum. Pale;” should read: “Mentum. Pale, occasionally with somewhat darker pigmentation”.

Nilothauma aleta Roback, 1960

Figures 1A, B, 17B

Additional material. 1 male, slide-mounted: BRAZIL, São Paulo, São Luís do Paraitinga, PE Serra do Mar, Núcleo Santa Virgínia, trilha Poço do Pito, afluenta Paraibuna, 23°20'09"S, 45°08'46"W, 15.ix.2006, light trap, M.R. Spies & A.E. Sieglöch leg.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX without setose dorsal lobe(s); gonostylus stout; acrostichals absent; anal point wide, covering most setae along posterior margin of tergite IX; inferior volsella slender.

Re-description. Male imago (n = 1). Total length 3.58 mm. Wing length 2.00 mm. Total length/wing length 1.79. Wing length/length of profemur 2.25.

Colouration. Head, thorax and abdomen brown; legs pale, except for ring of brown pigmentation in distal 1/2 to 2/3 of fore- and hind femora, in distal 1/3 of foretibia, in basal 1/8 of mid- and hind tibiae and in distal 1/3 of each tarsomere. Wing membrane apparently hyaline, but faint brown markings are visible when dark-field filter is applied.

Antenna. AR = 0.27. Thirteenth flagellomere 197 µm long.

Head. Temporal setae 7 in single row. Clypeus with 25 setae. Tentorium 123 µm long, maximum width 25 µm. Stipes not measurable. Palp segment lengths (in µm): 39, 34, 123, 147, 191. Third palpomere with 2 sensilla clavata subapically, longest 20 µm long. Fifth palpomere/third palpomere 1.55.




Figure 1. *Nilothauma aleta* Roback, 1960, adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Thorax. Dorsocentrals 12 in single row, acrostichals absent, prealars 3. Scutellum with 6 setae.

Wing. VR = 1.50. Brachiolum with 1 seta, R with 13 setae, R_1 with 18 setae, R_{4+5} with 21 setae, remaining veins bare.

Legs. Spur of fore tibia 44 μm long including 15 μm long scale. Mid-tibia with 1 spur, 15 μm long; hind tibia with 2 spurs, 25 and 29 μm long. Combs of both mid- and hind tibia 20 μm long. Width at apex of fore-tibia 39 μm , of mid-tibia 34 μm , of hind tibia 44 μm . Lengths and proportions of legs as in Table 1.

Hypopygium (Fig. 1A, B). Tergite IX without lobes, tapering to apex, with 22 short setae underneath anal point. Anal point lanceolate, 50 μm long, maximum width 37 μm . Tergite bands well developed. Laterosternite IX without setae. Phallapodeme 70 μm long; transverse sternapodeme 55 μm long. Gonocoxite 134 μm long. Inferior volsella straight, 52 μm long, 7 μm wide medially, with microtrichia and 8 simple setae apically. Superior volsella pediform, 17 μm long, 7 μm wide at base, covered with microtrichia and with 2 setae apically. Median volsella 7 μm long, with 2 simple setae, longest 12 μm . Gonostylus 95 μm long, straight. HR = 1.42. HV = 3.77.

Female adult and immatures. Unknown.

Remarks. Roback (1960) described *Nilothauma aleta* Roback, 1960 and *N. duena* Roback, 1960 from the Peruvian Amazon. In their revision of *Nilothauma*, Adam & Sæther, (1999) regarded the two species as not belonging to *Nilothauma* since they lack any projections on tergite IX and stated that they appear to belong in *Paratendipes* Kieffer. Later, Mendes and Andersen (2009) placed *Neelamia* Sponis and *Paranilothauma* Sponis as synonyms of *Nilothauma* and several new Neotropical species have been described demonstrating the large morphological variation in the genus. Mendes

Table 1. Lengths (in μm) and proportions of leg segments in *Nilothauma aleta* Roback, 1960, adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	887	601	837	522	404	256
P ₂	906	690	414	227	177	118
P ₃	1034	985	699	355	305	197
	ta ₅	LR	BV	SV	BR	
P ₁	148	1.39	1.74	1.77	2.3	
P ₂	89	0.60	3.29	2.30	2.3	
P ₃	108	0.71	2.56	2.89	5.0	

and Andersen (2009) emended the diagnosis of *Nilothauma* and both *N. aleta* and *N. duena* fit well into this diagnosis.

Distribution (Fig. 17B). The species was originally described from the Department of Huánuco, in the Peruvian Amazon by Roback (1960); the range is now extended to Serra do Mar (São Paulo State), in the Brazilian Atlantic Forest.

Nilothauma amazonense Mendes & Andersen, 2009

Figure 17A

Additional material. 1 male, slide-mounted: BRAZIL, Santa Catarina, Grão Pará, Cachoeira do Amado, #27, 28°08'57"S, 49°21'17"W, 16.xi.2012–08.i.2013, Malaise trap, L.C. Pinho, M.C. Novaes & M.F. Haddad leg. 1 male, slide-mounted: BRAZIL, Santa Catarina, São Francisco do Sul, Distrito do Saí, 26°11'42"S, 48°43'53"W, 18.i–18.iii.2020, Malaise trap #150, small stream, L.C. Pinho et al. leg.

Remarks. *Nilothauma amazonense* Mendes & Andersen, 2009 was described, based on a single male from the Amazon. The specimens of *N. amazonense* from southern Brazil, however, differ slightly from the holotype. Mendes and Andersen (2009) stated that hind ta₂ being shorter than ta₃ (ratio of ta₂/ta₃ length = 0.73) is one of the diagnostic characters of the species. However, in the specimens from southern Brazil, hind ta₂ and ta₃ are subequal in length (ratio of ta₂/ta₃ length = 0.94–0.97). Body size, measured as Total Length (TL) is also larger (TL of holotype = 1.53 mm; TL of southern populations = 2.00–2.05 mm). Similar differences in body size between Amazonian and southern Atlantic Forest populations were also found in *Beardius urupeatan* Pinho, Mendes & Andersen, 2009 [TL Amazon = 2.32–2.51, 2.38 (6); TL southern Atlantic Forest = 2.68–3.00, 2.96 (8)]. This intraspecific variation might be due to the higher temperature in the Amazon Region when compared to the localities in the southern parts of the Atlantic Forest. Populations of chironomid species inhabiting different habitats may show variation in voltinism and more rapid growth can result in smaller body size (Tokeshi 1995; Pinho et al. 2009).

Distribution (Fig. 17A). The species was originally described from the Amazonian Region by Mendes and Andersen (2009); the range is now extended to Santa Catarina State in southern Brazil.

***Nilothauma anamariae* Dantas & Hamada, 2017**

Figure 17D

Additional material. 1 male, slide-mounted: BRAZIL, Rondônia, Candeias do Jamari, Rio Preto, Ponte de Madeira, #01, 08°52'40"S, 63°38'02"W, 19–20.vii.2012, light trap, R. Boldrini & A.S. Fernandes leg. 1 male, slide mounted: BRAZIL, Mato Grosso, Ribeirão Cascalheira, Fazenda Campina Grande, Rio Suiá Miçu, 28–30.xi.2006, light trap, A.R. Calor, F.R. Silva & S. Mateus leg. 2 males, slide-mounted: BRAZIL, Mato Grosso, Ribeirão Cascalheira, Fazenda Campina Verde, Rio Suiá Miçu, 12°48.591'S, 52°06.925'W, 10.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg. 1 male, slide-mounted: BRAZIL, Pará State, Rurópolis, Rio Tambor, 29.x.2007, light trap, N. Hamada et al. leg.

Remarks. The inferior volsella can have up to 3–4 simple, curved setae apically.

Distribution (Fig. 17D). The species was originally described from the Rio Grande do Sul State in southern Brazil by Dantas and Hamada (2017); the range is now extended to Mato Grosso, Rondônia and Pará States in central and northern Brazil.

***Nilothauma aripuanense* Mendes & Andersen, 2009**

Figure 17B

Additional material. 2 males, slide-mounted: BRAZIL, Rondônia, Candeias do Jamari, Rio Preto, Ponte de Madeira, #01, 08°52'40"S, 63°38'02"W, 19–20.vii.2012, light trap, R. Boldrini & A.S. Fernandes leg. 2 males, slide-mounted: BRAZIL, Rondônia, Teixeiraópolis, Balneário com Cachoeira, 10°55'20"S, 62°22'34"W, 03.ix.2012, light #13, N. Hamada, R. Boldrini, A.S. Fernandes & J.M. Cavalcante leg. 1 male, slide-mounted: BRAZIL, Roraima, Boa Vista, Rio Cauamé, 02°52'06"N, 60°44'24"W, 9.iii.2009, light trap, L.M. Fusari leg. 1 male, slide-mounted: BRAZIL, Roraima, Boa Vista, BR-174, Igarapé Água Boa, 02°43'32"N, 60°48'43"W, 2014, N. Hamada leg. 1 male, slide-mounted: BRAZIL, Amazonas, Presidente Figueiredo, AM-240 Km 20, Balneário Sossego da Panteira, Igarapé da Onça, 02°02'31"S, 59°51'05"W, 02.vii.2008, light trap, C. Azevedo leg. 1 male, slide-mounted: BRAZIL, Amazonas, upper Rio Marauaiá, downstream of Cachoeira Santo Antônio, surface float skimmed, 22.i.1963, E.J. Fittkau leg. (A485, ZSM).

Distribution (Fig. 17B). The species was originally described from the Amazonas and Mato Grosso States by Mendes and Andersen (2009); the range is now extended to Roraima and Rondônia States in the Brazilian Amazon.

***Nilothauma calori* Mendes & Andersen, 2009**

Figures 2A, 16C

Additional material. 1 male, slide-mounted: BRAZIL, Amazonas, Manaus, Reserva Florestal Adolfo Ducke, Igarapé Bolívia, 02°49'15"S, 59°56'31"W, 9–12.xi.2008,




Figure 2. Variation in superior volsella **A** dorsolateral view of superior volsella of *Nilothauma calori* Mendes & Andersen, 2009 **B** lateral view of superior volsella of *Nilothauma complicatum* Mendes & Andersen, 2009.

Malaise trap suspensa 2, N. Hamada et al. leg. 2 males, slide-mounted: BRAZIL, Mato Grosso, Cuiabá, 10–11.x.1965, Brundin net, E.J. Fittkau leg. (A 580, ZSM).

Remarks. Mendes and Andersen (2009) stated that the superior volsella has a “lateral strongly sclerotized, spine-like projection”. In dorsolateral view (Fig. 2A), it can be seen that this spine-like projection originates from the base of the volsella and is equally long as the volsella proper.

Distribution (Fig. 16C). The species was originally described from Acre State by Mendes and Andersen (2009); the range is now extended to the Mato Grosso and Amazonas States.

Nilothauma complicatum Mendes & Andersen, 2009

Figures 2B, 16A

Additional material. 1 male, slide-mounted: BRAZIL, Rondônia, Candeias do Jamari, Rio Preto, Ponte de Madeira, #01, 08°52'40"S, 63°38'02"W, 19–20.vii.2012, light trap, R. Boldrini & A.S. Fernandes leg. 3 males, slide-mounted: BRAZIL, Mato Grosso, Nova Xavantina, Fazenda Sr. Queté, Córrego Voadeira, 14°32.187'S, 52°30.902'W, 16.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg. 2 males, slide-mounted, as previous, except: Córrego Cachoeira, 14°32.817'S, 52°31.395'W. 1 male, slide-mounted, as previous, except: 14°41.577'S, 52°27.203'W, 13.x.2007. 2 males, slide-mounted, as previous, except: Estrada p/ Rancho Helena, Córrego Ponte de

Pedra, 14°47.908'S, 52°37.226'W, 15.x.2007. 1 male, slide-mounted, as previous, except: Córrego Voadeira, 14°41.577'S, 52°27.203'W, 13.x.2007. 1 male, slide-mounted: BRAZIL, Mato Grosso, Ribeirão Cascalheira, Estrada Fazenda Manaus, 1° afluente Rio Bonito, 12°57.088'S, 51°52.480'W, 08.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg.

Remarks. Mendes and Andersen (2009) stated that the superior volsella has a “marginal row of flattened setae”. In lateral view (Fig. 2B), it can be seen that the volsella is quite wide medially with a row of lamellae apparently covered with weak microtrichia.

Distribution (Fig. 16A). The species was originally described by Mendes and Andersen (2009), based on a single male from the Espírito Santo State; the range is now extended to the Mato Grosso and Rondônia States in central and northern Brazil.

Nilothauma duena Roback, 1960

Figures 3A, B, 16C

Additional material. 1 male, slide-mounted: BRAZIL, Bahia, Camacan, Fazenda do Waldemar da farmácia, Córrego abaixo da represa de abastecimento, 15°25'16"S, 39°33'57"W, 300 m a.s.l., 05.viii.2008, light trap, A.R. Calor, L.S. Lecci, L.C. Pinho & R.A. Moretto leg. 1 male, slide-mounted: BRAZIL, São Paulo, Pindamonhangaba, Fazenda São Sebastião, Afluente Cedro 3, 22°50'16"S, 45°28'27"W, 18.ix.2006, light trap, M.R. Spies & A.E. Sieglösch leg.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX without setose dorsal lobe(s); anal point slightly spatulate; wing unmarked; abdominal tergites I–VIII with basal half light brown, distal half pale; gonostylus and inferior volsella stout.

Re-description. Male imago (n = 2, unless otherwise stated). Total length 2.37–2.76 mm. Wing length 1.56–1.68 mm. Total length/wing length 1.58–1.66. Wing length/length of profemur 2.60–2.62.

Colouration. Head and thorax light brown; legs pale except for brown pigmentation in basal 1/3 and distal 1/5 of fore femur, basal 1/5 and distal 1/3 of fore tibia, distal 1/8 of mid- and hind femora, basal 1/8 and distal 1/8 of mid- and hind tibiae and entire tarsi. Abdominal tergites I–VIII with basal half light brown, distal half pale; hypopygium light brown. Wing membrane hyaline.

Antenna. AR = 0.17–0.18. Thirteenth flagellomere 108–118 μ m long.

Head. Temporal setae 9–10 in single row. Clypeus with 13–15 setae. Tentorium 98 (1) μ m long, maximum width 20 (1) μ m. Stipes not measurable. Palp segment lengths (in μ m): 17–25, 17–25, 32 (1), 35 (1), 62 (1). Third palpomere with 2 sensilla clavata subapically, longest 15 μ m long. Fifth palpomere/third palpomere 1.92 (1).

Thorax. Dorsocentrals 16–17 in single row, acrostichals 14–16, prealars 3. Scutellum with 2–3 setae.

Wing. VR = 1.55–1.56. Brachiolum with 1 seta, R with 12–13 setae, R₁ with 16–17, R₄₊₅ with 5–22 setae, remaining veins bare.




Figure 3. *Nilothauma duena* Roback, 1960, adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Legs. Spur of fore tibia 54–59 μm long including 15–20 μm long scale. Mid-tibia with 1 spur, 25–29 μm long; hind tibia with 2 spurs, 49–51 and 28–31 μm long. Combs of mid-tibia 15–20 μm long, of hind tibia 18–25 μm long. Width at apex of fore tibia 48–50 μm , of mid-tibia 48–50 μm , of hind tibia 49–59 μm . Lengths and proportions of legs as in Table 2.

Hypopygium (Fig. 3A, B). Tergite IX without dorsal lobes, with triangular posterior margin with 13–15 weak setae along posterior margin to each side of base of anal point. Anal point spatulate, 22–30 μm long, maximum width 10–12 μm . Tergite bands well developed. Laterosternite IX with 1–2 setae. Phallapodeme 47–60 μm long; transverse sternapodeme 55–62 μm long. Gonocoxite 112–125 μm long. Inferior volsella straight, 40–45 μm long, 15–20 μm wide medially, with microtrichia and 10–11 strong, simple setae apically. Superior volsella tapering to apex, 20–37 μm long, 12–14 μm wide at base, covered with microtrichia and apparently bare at tip. Median volsella 7–10 μm long, with 3–4 setae, longest 20–22 μm long. Gonostylus 87–90 μm long, straight. HR = 1.29–1.43. HV = 2.72–3.17.

Female adult and immatures. Unknown.

Remarks. See remarks for *N. aleta* Roback, 1960.

Distribution (Fig. 16C). The species was originally described by Roback (1960) from the Department of Huánuco, in the Peruvian Amazon; the range is now extended to Serra Bonita (Bahia State) and Serra do Mar (São Paulo State), in the Brazilian Atlantic Forest.

Table 2. Lengths (in μm) and proportions of leg segments in *Nilothauma duena* Roback, 1960, adult males ($n = 2$).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	601–640	453–502	660–690	374–376	276–278	207–217
P ₂	621–670	453–473	296–305	148–158	108–110	69–79
P ₃	739–778	670–699	404–443	217–236	207–210	148–150
	ta ₅	LR	BV	SV	BR	
P ₁	105–108	1.37–1.46	1.72–1.74	1.60–1.66	1.8–2.7	
P ₂	49–59	0.65–0.66	3.48–3.49	3.56–3.74	1.8–2.7	
P ₃	79–99	0.60–0.63	2.78–2.79	3.33–3.49	3.6–4.7	

***Nilothauma fittkaui* (Soponis, 1987)**

Figure 17A

Additional material. 2 males, slide-mounted: BRAZIL, São Paulo, Campos do Jordão, PE Campos do Jordão, Córrego Canhambora, 1538 m a.s.l., 22°41'44"S, 45°29'30"W, 13.i.2006, light trap, M.R. Spies leg. 1 male, slide-mounted, COSTA RICA, La Selva, 03.iv.1993, Malaise trap, O.A. Sæther leg.

Distribution (Fig. 17A). The species was described by Soponis (1987) from Amazonas and later recorded from Acre, Espírito Santo and Para States in Brazil and from Ecuador by Mendes and Andersen (2009). The range is now extended south to São Paulo State in Brazil and north to Costa Rica in Central America.

***Nilothauma hamadae* sp. nov.**

<http://zoobank.org/40ABE99F-793E-4688-9F2E-4D8B21B15E37>

Figures 4A, B, 16A

Type material. *Holotype* male, slide-mounted: BRAZIL, Amazonas, Barcelos, Rio Aracá, Foz do Igarapé Cuieiras, 00°19'15"N, 63°16'15"W, 35 m a.s.l., 30.vii–01.viii.2009, light trap #11, N. Hamada et al. leg. (UFSC).

Etymology. The specific epithet is a noun in the genitive case which honours Neusa Hamada for her great contribution to the knowledge of Amazonian Chironomidae.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX with single, median setose dorsal lobe, consisting of a low, but wide protuberance with about 25 strong setae; anal point very broad (about half the width of tergite IX) and lanceolate; median volsella separated from superior volsella.

Description. Male imago (n = 1). Total length 1.94 mm. Wing length 1.13 mm. Total length/wing length 1.71. Wing length/length of profemur 2.30.

Colouration. Head, thorax and abdomen medium brown. Legs light brown. Wing membrane hyaline.

Antenna. AR = 0.40. Thirteenth flagellomere 217 μm long.

Head. Temporal setae 7 in single row. Clypeus with 15 setae. Tentorium 75 μm long, maximum width 12 μm . Stipes 92 μm long. Palp segment lengths (in μm): 25,




Figure 4. *Nilothauma hamadae* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

25, 75, 127, 144. Third palpomere with 2 sensilla clavata subapically, longest 20 μm long. Fifth palpomere/third palpomere 1.93.

Thorax. Dorsocentrals 7 in single row, acrostichals 10, prealars 2. Scutellum with 4 setae.

Wing. VR = 1.37. Brachiolum with 1 seta, R with 13 setae, R_1 with 7 setae, R_{4+5} with 14 setae, remaining veins bare.

Legs. Spur of fore tibia 34 μm long including 12 μm long scale. Mid-tibia with 1 spur, 15 μm long; hind tibia with 2 spurs, 20 and 25 μm long. Combs of mid-tibia 12 μm long, of hind tibia 15 μm long. Width at apex of fore tibia 34 μm , of mid-tibia 34 μm , of hind tibia 39 μm . Lengths and proportions of legs as in Table 3.

Hypopygium (Fig. 4A, B). Tergite IX narrow, tapering to apex, with central rounded lobe bearing 25 simple, strong setae; with 9 simple setae along posterior margin underneath the anal point. Anal point lanceolate, 42 μm long, 27 μm wide. Tergite bands lacking. Laterosternite IX with 1 seta. Phallapodeme 40 μm long; transverse sternapodeme 15 μm long. Gonocoxite 75 μm long. Inferior volsella strongly curved,

Table 3. Lengths (in μm) and proportions of leg segments in *Nilothauma hamadae* sp. nov., adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	493	364	–	–	–	–
P ₂	473	364	217	108	79	49
P ₃	532	542	315	167	158	99
	ta ₅	LR	BV	SV	BR	
P ₁	–	–	–	–	–	–
P ₂	49	0.59	3.62	3.86	3.3	–
P ₃	69	–	–	–	5.0	–

37 μm long, 7 μm wide medially, with microtrichia in basal half, with 4 simple setae apically and 1 simple seta subapically. Superior volsella 30 μm long, 4 μm wide at base, covered with microtrichia and fringed at apex. Median volsella 7 μm long, with 2 simple setae, longest 10 μm . Gonostylus 112 μm long, with basal half strongly curved and distal half straight. HR = 0.67. HV = 1.73.

Female adult and immatures. Unknown.

Distribution (Fig. 16A). Known from Barcelos (Amazonas State), in the Brazilian Amazon.

Nilothauma jaraguaense Mendes & Andersen, 2009

Figure 17D

Additional material. 1 male, slide-mounted: BRAZIL, São Paulo, Salesópolis, EB Boraceia, Rio Claro, Poço Verde, 18.ix.2002, light trap, A.S. Melo, C.G. Froehlich, R. Mariano, A. Prather & R. Blahnik leg. 1 male, slide-mounted: BRAZIL, São Paulo, Jundiá, PE Serra do Japi, 23.ix.2008, light trap, R. Mariano & L.S. Lecci leg.

Distribution (Fig. 17D). The species was described by Mendes and Andersen (2009), based on a single male from Parque Estadual do Jaraguá in São Paulo State, Brazil.

Nilothauma jupau sp. nov.

<http://zoobank.org/3F55ECB7-8CFC-4229-B33A-3FD9520A23B0>

Figures 5A–C, 16A

Type material. Holotype male, slide-mounted: BRAZIL, Rondônia, Teixeirópolis, Balneário com Cachoeira, 10°55'20"S, 62°22'34"W, 03.ix.2012, light trap #13, N. Hamada, R. Boldrini, A.S. Fernandes & J.M. Cavalcante leg. (UFSC). **Paratype:** 1 male, slide-mounted, same data as holotype (INPA).

Etymology. The specific epithet honours the Jupaú, indigenous people from Rondônia State, Brazilian Amazon. The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX with thorn and without setose dorsal lobe(s); anal point




Figure 5. *Nilothauma jupau* sp. nov. adult male **A** wing **B** hypopygium, dorsal view **C** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

spatulate; wing with conspicuous dark markings; abdominal tergites II, III, VI, VII and VIII dark brown.

Description. Male imago (n = 2, unless otherwise stated). Total length 2.04–2.14 mm. Wing length 1.07–1.14 mm. Total length/wing length 1.78–2.01. Wing length/length of profemur 2.03–2.27.

Colouration. Head and thorax brown; legs pale, except for entire fore femur, distal half of mid- and hind femora and distal 1/3 of fore- and hind tibiae with brown pigmentation; abdomen pale, except for brown pigmentation in segments II, III, VI, VII and VIII. Wing membrane with extensive dark markings.

Antenna. AR = 0.28–0.32. Thirteenth flagellomere 115–134 μ m long.

Table 4. Lengths (in μm) and proportions of leg segments in *Nilothauma jupau* sp. nov., adult males ($n = 2$).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	502–522	404–424	–	–	–	–
P ₂	483–522	335–345	180–187	87–89	57–59	37–39
P ₃	542–571	512–522	270–276	138–148	138–148	85–89
	ta ₃	LR	BV	SV	BR	
P ₁	–	–	–	–	–	
P ₂	28–30	0.54–0.56	3.22–3.31	4.22–4.56	1.8–2.8	
P ₃	59–69	0.53–0.54	2.93–3.23	3.82–3.96	1.5–3.2	

Head. Temporal setae 7–8 in single row. Clypeus with 20–24 setae. Tentorium 57–72 μm long, maximum width 15–17 μm . Stipes 95–100 μm long. Palp segment lengths (in μm): 13–14, 30–32, 85–95, 105–117, 115–134. Third palpomere with 2–3 sensilla clavata subapically, longest 14–15 μm . Fifth palpomere/third palpomere 1.21–1.69.

Thorax. Dorsocentrals 6–7 in single row, acrostichals 8–14, prealars 2. Scutellum with 2 setae.

Wing (Fig. 5A). VR = 1.63–1.64. Brachiolum with 1 seta, R with 8–10 setae, R₁ with 6 setae, R₄₊₅ with 11–15 setae, remaining veins bare.

Legs. Spur of fore tibia 39–49 μm long including 20–25 μm long scale. Mid-tibia with 1 spur, 20–25 μm long; hind tibia with 2 spurs, 20–25 and 23–28 μm long. Combs of mid-tibia 15–20 μm long, of hind tibia 15–20 μm long. Width at apex of fore tibia 44 μm , of mid-tibia 39–44 μm , of hind tibia 47–51 μm . Lengths and proportions of legs as in Table 4.

Hypopygium (Fig. 5B, C). Tergite IX without dorsal lobes, with rounded posterior margin, with 4–5 clustered setae anteriorly to base of anal point and 8–12 weaker setae to each side of anal point. Anal point spatulate, 40–42 μm long, maximum width 7–10 μm . Tergite bands well developed. Laterosternite IX with 2–3 setae, with thorn. Phallapodeme 42–52 μm long; transverse sternapodeme 20–22 μm long. Gonocoxite 70–75 μm long, with longer microtrichia dorsomedially. Inferior volsella slightly curved, 30–32 μm long, 15–18 μm wide medially, with microtrichia and 5–6 simple setae in apical one third. Superior volsella slender, 12–20 μm long, 4–5 μm wide at base, covered with microtrichia and with 2 setae at apex, longest 8–13 μm . Median volsella consisting of small tubercle situated underneath superior volsella, 6–7 μm long, with 2 setae at apex, longest 6–7 μm long. Gonostylus 70–92 μm long, straight. HR = 0.81–1.00. HV = 2.21–3.05.

Female adult and immatures. Unknown.

Distribution (Fig. 16A). Known from Rondônia State, Brazilian Amazon.

***Nilothauma karitiana* sp. nov.**

<http://zoobank.org/20FE2D58-13AD-47D5-9414-9EB6B7BFC321>

Figures 6A–C, 16C

Type material. *Holotype* male, slide-mounted: BRAZIL, Rondônia, Candeias do Jamari, Rio Preto, Ponte de Madeira, 08°52'40"S, 63°38'02"W, 19–20.vii.2012, light




Figure 6. *Nilothauma karitiana* sp. nov. adult male **A** hypopygium, dorsal view **B** superior volsella, dorsal view **C** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

trap #01, R. Boldrini & A.S. Fernandes leg. (UFSC). **Paratype:** 1 male adult, slide-mounted: BRAZIL, Amazonas, Barcelos, Rio Aracá, #9, 69 m a.s.l., 00°24'39"N, 63°23'12"W, 28.vii–06.viii.2009, light trap #3, N. Hamada et al. leg. (INPA).

Etymology. The specific epithet honours the Karitiana, indigenous people from the Rio Jamari Basin in the Rondônia State (Brazil). The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX without setose dorsal lobe(s); anal point spatulate; wing unmarked; superior volsella fused to median volsella; gonostylus very long, narrow basally and apically, swollen medially.

Table 5. Lengths (in μm) and proportions of leg segments in *Nilothauma karitiana* sp. nov., adult males ($n = 1-2$).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	847–887	729–808	1054	542	424	335
P ₂	739–798	532–561	374	158	108	69
P ₃	896–965	867–926	493–522	246–256	240–246	144–148
	ta ₅	LR	BV	SV	BR	
P ₁	138	1.30	1.91	1.61	2.2	
P ₂	49	0.67	4.51	3.63	3.2	
P ₃	95–99	0.56–0.57	3.01–3.27	3.58–3.62	4.0–5.0	

Description. Male imago ($n = 2$, unless otherwise stated). Total length 3.32–3.78 mm. Wing length 1.71–1.83 mm. Total length/wing length 1.94–2.07. Wing length/length of profemur 2.02–2.06.

Colouration. Head, thorax, legs and abdomen uniformly light brown. Wing membrane hyaline.

Antenna. AR = 0.22 (1). Thirteenth flagellomere 217 μm long.

Head. Temporal setae 9 (1) in single row. Clypeus with 10–11 setae. Tentorium 105 (1) μm long, maximum width 17 (1) μm . Stipes 122 (1) μm long. Palp segment lengths (in μm): 25–35, 27–37, 60 (1), 80 (1), 154 (1). Third palpomere with 3 (1) sensilla clavata subapically, longest 20 (1) μm long. Fifth palpomere/third palpomere 2.57

Thorax. Dorsocentrals 9–13 in single row, acrostichals 12, prealars 2–3. Scutellum with 2 setae.

Wing. VR = 1.33–1.46. Brachiolum with 1 seta, R with 11–14 setae, R₁ with 10–12 setae, R₄₊₅ with 3–4 setae at apex, remaining veins bare.

Legs. Spur of fore tibia 59–64 μm long including 17–20 μm long scale. Mid-tibia with 1 spur, 23–25 μm long; hind tibia with 2 spurs, 23–25 and 29–33 μm long. Combs of mid-tibia 15–20 μm long, of hind tibia 20–23 μm long. Width at apex of fore tibia 59 μm , of mid-tibia 59 μm , of hind tibia 64 μm . Lengths and proportions of legs as in Table 5.

Hypopygium (Fig. 6A, B). Tergite IX without dorsal lobes, tapering to apex, with 2–3 median and 8–9 setae to each side of anal point. Anal point spatulate, 55–57 μm long, 17–20 μm wide. Tergite bands well developed. Laterosternite IX with 1–2 setae. Phallapodeme 82–90 μm long; transverse sternapodeme 42–55 μm long. Gonocoxite 138–142 μm long. Inferior volsella slightly curved, 97–107 μm long, 17–20 μm wide medially, with microtrichia and 6–7 simple setae subapically. Superior volsella digitiform, 55–65 μm long, 17–20 μm wide at base, covered with microtrichia and with 4 setae apically. Median volsella fused to superior volsella, consisting of 2–4 small tubercles each bearing single, simple seta, longest 22–25 μm . Gonostylus 204–232 μm long, straight, narrow basally and apically, swollen medially. HR = 0.62–0.70. HV = 1.62–1.63.

Female adult and immatures. Unknown.

Distribution (Fig. 16C). Known from Rondônia and Amazonas States, in the Brazilian Amazon.

***Nilothauma leccii* sp. nov.**

<http://zoobank.org/CBFC7D88-17AB-4BF8-9CBA-2995FDAB798B>

Figures 7A, B, 16B

Type material. *Holotype* male, slide-mounted: BRAZIL, São Paulo, São Sebastião, Rio das Pedras, 23°44'27"S, 45°37'12"W, 28.x.2005, light trap, A.R. Calor et al. leg. (UFSC).

Etymology. The specific epithet is a noun in the genitive case honouring Lucas Silveira Lecci, for his friendship and prolific fieldwork.

Diagnostic characters. The male can be separated from its congeners by its large size combined with unmarked wing; spatulate anal point; superior volsella leaf-shaped; inferior volsella with strong, split setae and digitiform and strongly setose gonostylus.

Description. Male imago (n = 1). Total length 4.35 mm. Wing length 1.96 mm. Total length/wing length 2.22. Wing length/length of profemur 2.45.

Colouration. Thorax and legs brown, abdomen light brown. Wing membrane without dark markings.

Antenna. AR = 0.16. Thirteenth flagellomere 124 µm long.

Head. Temporal setae 6 in single row. Clypeus with 15 setae. Tentorium 113 µm long, maximum width 25 µm. Stipes not measurable. Palp segment lengths (in µm): 37, 33, 74, 107, 138. Third palpomere with 2 sensilla clavata subapically, longest about 25 µm long. Fifth palpomere/third palpomere 1.86.

Thorax. Antepronotum with 4 setae. Dorsocentrals 17 partly biserial posterior, acrostichals 14, prealars 6. Scutellum with 13 setae.

Wing. VR = 1.53. Brachiolum with 2 setae, R with 15 setae, R₁ with 24 setae, R₄₊₅ with 31 setae, remaining veins bare.

Legs. Spur of fore tibia 65 µm long including 41 µm long scale. Mid-tibia with 1 spur, 47 µm long; hind tibia with 2 spurs, 43 and 65 µm long. Combs of mid-tibia 29 µm long, of hind tibia 47 µm long. Width at apex of fore tibia 69 µm, of mid-tibia 73 µm, of hind tibia 89 µm. Lengths and proportions of legs as in Table 6.

Hypopygium (Fig. 7A, B). Tergite IX without dorsal lobes, posterior margin subtriangular with 7 setae to each side of the anal point. Anal point spatulate, 59 µm long, 10 µm wide basally, 17 µm wide medially. Tergite bands not continuous. Laterosternite IX with 4 setae. Phallapodeme 61 µm long; transverse sternapodeme 35 µm long. Gonocoxite 171 µm long. Inferior volsella weakly curved, 113 µm long, 15 µm wide subapically, with microtrichia and 7 strong, apically split setae. Superior volsella leaf-shaped, 69 µm long, 10 µm wide at base, 23 µm wide medially, covered with

Table 6. Lengths (in µm) and proportions of leg segments in *Nilothauma leccii* sp. nov., adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	801	670	972	490	384	286
P ₂	874	605	400	180	139	90
P ₃	964	989	596	302	261	155
	ta ₅	LR	BV	SV	BR	
P ₁	139	1.45	1.88	1.51	1.7	
P ₂	65	0.66	3.97	3.69	4.3	
P ₃	98	0.60	3.09	3.27	4.8	




Figure 7. *Nilothauma leccii* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

microtrichia and with few weak setae ventrally and along inner margin. Median volsella narrow, 48 μm long, covered with microtrichia and with 4 setae on tubercles at apex, setae about 23 μm long. Gonostylus digitiform, strongly setose, 163 μm long, 37 μm wide medially. HR = 1.05. HV = 2.67.

Female adult and immatures. Unknown.

Distribution (Fig. 16B). Only known from São Paulo State in Brazil.

***Nilothauma mariano* sp. nov.**

<http://zoobank.org/2F9585A3-AE43-47C7-BD91-C4EE9B23EEBC>

Figures 8A, B, 16B

Type material. *Holotype* male, slide-mounted: BRAZIL, Bahia, Barreiras, Rio de Janeiro, cachoeira Acaba Vidas, 11°53'40"S, 45°36'57"W, 722 m a.s.l., 14.x.2008, light trap, A.R. Calor, R. Mariano & S. Mateus leg. (UFSC).




Figure 8. *Nilothauma marianoi* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Etymology. The specific epithet is a noun in the genitive case honouring Rodolfo Mariano, for his friendship and prolific fieldwork.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: wing without dark markings; tergite IX without setose dorsal lobe(s) or thorns, with single, strong median seta, with narrowly triangular posterior margin and small, apical, parallel-sided anal point.

Description. Male imago (n = 1). Total length 3.51 mm. Wing length 1.45 mm. Total length/wing length 2.43. Wing length/length of profemur 2.27.

Colouration. Thorax and legs brown, abdomen light brown. Wing membrane without dark markings.

Antenna. AR = 0.19. Thirteenth flagellomere 152 μ m long.

Head. Temporal setae 6 in partly double row. Clypeus with 9 setae. Tentorium 98 μ m long, maximum width 18 μ m. Stipes not measurable. Palp segment lengths (in μ m): 17, 18, 44, 99, 117. Sensilla clavata on third palpomere not discernable. Fifth palpomere/third palpomere 2.66.

Thorax. Dorsocentrals 6 in single row, acrostichals 4, prealars 2. Scutellum with 2 setae.

Table 7. Lengths (in μm) and proportions of leg segments in *Nilothauma mariano* sp. nov., adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	637	458	645	401	310	221
P ₂	621	425	261	131	90	65
P ₃	694	686	384	180	180	114
	ta ₃	LR	BV	SV	BR	
P ₁	106	1.41	1.68	1.70	2.6	
P ₂	49	0.62	3.90	4.05	5.0	
P ₃	74	0.55	3.22	3.60	7.1	

Wing. VR = 1.39. Brachiolum with 1 seta, R with 8 setae, R₄₊₅ with 1 apical seta, remaining veins bare.

Legs. Spur of fore tibia 68 μm long including 28 μm long scale. Mid-tibia with 1 spur, 47 μm long; hind tibia with 2 spurs, 43 and 61 μm long. Combs of mid-tibia 21 μm long, of hind tibia 26 μm long. Width at apex of fore tibia 47 μm , of mid-tibia 48 μm , of hind tibia 52 μm . Lengths and proportions of legs as in Table 7.

Hypopygium (Fig. 8A, B). Tergite IX without dorsal lobes, with single, median, strong setae, posterior margin narrowly subtriangular with 7 setae to each side. Anal point situated apically, small, parallel-sided with rounded apex, 14 μm long, 4 μm wide basally, 3 μm wide medially. Tergite bands lacking. Laterosternite IX with 2 setae. Phallapodeme 51 μm long; transverse sternapodeme 11 μm long. Gonocoxite 104 μm long. Inferior volsella weakly curved, 103 μm long, 11 μm wide subapically, with microtrichia and 18 setae in apical one-half. Superior volsella subquadrangular, 48 μm long, 17 μm wide medially, covered with microtrichia. Median volsella consisting of 14 μm long tubercle, covered with microtrichia and with 1 strong apical seta, setae about 19 μm long. Gonostylus nearly straight, 104 μm long, 10 μm wide medially, 17 μm wide subapically. HR = 0.72. HV = 2.44.

Female adult and immatures. Unknown.

Distribution (Fig. 16B). Only known from Bahia State in Brazil.

Nilothauma mateusi sp. nov.

<http://zoobank.org/679EA037-9083-4F6D-9460-3F9F8E590EA2>

Figures 9A, B; 16B

Type material. Holotype male, slide-mounted: BRAZIL, Mato Grosso, Nova Xavantina, Fazenda Sr. Queté, Córrego Cachoeira, 14°32.817'S, 52°31.395'W, 16.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg. (UFSC).

Etymology. The specific epithet is a noun in the genitive case honouring Sidnei Mateus, for his friendship and prolific fieldwork.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: wing without markings; tergite IX with pair of rounded lobes submedially with about 14 long setae; anal point parallel-sided; superior volsella small, subtriangular, projecting medially, with 2 setae on tubercles apically.




Figure 9. *Nilothauma mateusi* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Description. Male imago (n = 1). Total length 2.09 mm. Wing length 0.87 mm. Total length/wing length 2.40. Wing length/length of profemur 2.41.

Colouration. Thorax and legs brown, abdomen light brown. Wing membrane without dark markings.

Antenna. AR = 0.19. Thirteenth flagellomere 82 μ m long.

Head. Temporal setae 7 in single row. Clypeus with 9 setae. Tentorium 55 μ m long, maximum width 12 μ m. Stipes not measurable. Palp segment I–III lengths (in μ m): 21, 19, 55; remaining palp segments lost. Third palpomere with 2 sensilla clavata subapically, longest about 10 μ m.

Thorax. Dorsocentrals 8 in single row, acrostichals 6, prealars 2. Scutellum with 2 setae.

Wing. VR = 1.64. Brachiolum with 1 seta, R with 6 setae, R_1 with 2 setae, R_{4+5} with 4 setae, remaining veins bare.

Legs. Spur of fore tibia 37 μ m long including 18 μ m long scale. Mid-tibia with 1 spur, 23 μ m long; hind tibia with 2 spurs, 19 and 28 μ m long. Combs of mid-tibia 14 μ m long, of hind tibia 17 μ m long. Width at apex of fore tibia 29 μ m, of mid-tibia 30 μ m, of hind tibia 33 μ m. Lengths and proportions of legs as in Table 8.

Table 8. Lengths (in μm) and proportions of leg segments in *Nilothauma mateusi* sp. nov., adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	360	272	340	136	112	68
P ₂	352	248	140	64	52	32
P ₃	416	392	208	108	112	72
	ta ₅	LR	BV	SV	BR	
P ₁	40	1.25	2.73	1.86	2.0	
P ₂	24	0.56	4.30	4.29	2.8	
P ₃	40	0.53	3.06	3.88	4.6	

Hypopygium (Fig. 9A, B). Tergite IX with 4 weak setae to each side of the anal point and pair of rounded lobes submedially, each with about 14 long setae, longest about 50 μm long. Anal point parallel-sided with rounded apex, 23 μm long, 12 μm wide basally, 8 μm wide medially. Tergite bands not continuous. Laterosternite IX with 1 seta. Phallapodeme 35 μm long; transverse sternapodeme 19 μm long. Gonocoxite 73 μm long. Inferior volsella straight, 43 μm long, 7 μm wide subapically, with microtrichia and 4 strong apical setae. Superior volsella projecting medially, subtriangular with two apical tubercles, 14 μm long, 7 μm wide at base, 4 μm wide subapically, covered with microtrichia and with 2 apical setae, longest 9 μm long. Median volsella consisting of single strong tubercle, about 12 μm long, with single 10 μm long setae at apex. Gonostylus curved, 101 μm long, 10 μm wide medially, 14 μm wide subapically. HR = 0.72. HV = 2.07.

Female adult and immatures. Unknown.

Distribution (Fig. 16B). Only known from Mato Grosso State in Brazil.

***Nilothauma maya* sp. nov.**

<http://zoobank.org/0165F284-D9E4-4FF1-B207-C4EE505A0E00>

Figures 10A, B, 16B

Type material. Holotype male, slide-mounted: MEXICO, Campeche, Calacmul, Ejido Nuevo Becan, El Chorro, 18°35'26"N, 89°15'29"W, 130 m a.s.l., 30.iv.1997, light trap, A. Contreras-Ramos et al. leg. (ZMBN).

Etymology. Named after the Maya people, who used to live in the area. The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: pale brown species; wing without markings; tergite IX without setose dorsal lobe(s) or spine; anal point spatulate; superior volsella slender, curved, tapering; gonostylus curved, with strong setae on protruberance on inner margin in basal one-third.

Description. Male imago (n = 1). Total length 2.43 mm. Wing length 1.12 mm. Total length/wing length 2.17. Wing length/length of profemur 2.15.

Colouration. Pale brown. Wing membrane without dark markings.

Antenna. AR = 0.18. Thirteenth flagellomere 112 μm long.




Figure 10. *Nilothauma maya* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Head. Temporal setae 4 in single row. Clypeus with 13 setae. Tentorium and stipes not measurable. Palp segment lengths (in μm): 21, 28, 65, 117, 144. Third palpomere with 2 sensilla clavata subapically, longest 19 μm long. Fifth palpomere/third palpomere 2.22.

Thorax. Dorsocentrals 8 in single row, acrostichals 8, prealars 2. Scutellum with 2 setae.

Wing. VR = 1.44. Brachiolum with 1 seta, R with 10 setae, R_1 with 12 setae, R_{4+5} with 15 setae, remaining veins bare.

Legs. Spur of fore tibia 48 μm long including 21 μm long scale. Mid-tibia with 1 spur, 26 μm long; hind tibia with 2 spurs, 25 and 33 μm long. Combs of mid-tibia 12 μm long, of hind tibia 17 μm long. Width at apex of fore tibia 36 μm , of mid-tibia 40 μm , of hind tibia 44 μm . Lengths and proportions of legs as in Table 9.

Hypopygium (Fig. 10A, B). Tergite IX without lobes, with 8 setae above anal point and 5 somewhat weaker setae to each side of anal point. Anal point spatulate, 36 μm long, maximum width 11 μm . Tergite bands continuous. Laterosternite IX with 2 setae. Phallapodeme 48 μm long; transverse sternapodeme 27 μm long. Gonocoxite 104 μm long. Inferior volsella curved, 55 μm long, 14 μm wide subapically, with microtrichia and 4 strong apical setae. Superior volsella slender, curved, tapering, 54 μm

Table 9. Lengths (in μm) and proportions of leg segments in *Nilothauma maya* sp. nov., adult male (n = 1).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	520	372	–	–	–	–
P ₂	484	308	208	76	56	32
P ₃	504	484	232	132	140	88
	ta ₅	LR	BV	SV	BR	
P ₁	–	–	–	–	–	–
P ₂	28	0.68	5.21	3.81	3.4	
P ₃	48	0.48	2.99	4.26	4.7	

long, 11 μm wide at base, 3 μm wide subapically, covered with microtrichia and with 2 weak apical setae. Median volsella consisting of two small tubercles, about 4 μm long, each with single setae at apex, longest 15 μm long. Gonostylus 131 μm long, curved, with single, strong setae on protuberance on inner margin at 27 μm from base, setae 21 μm long. HR = 0.79. HV = 1.85.

Female adult and immatures. Unknown.

Remarks. The slide is distorted and the drawings are composites of left and right side.

Distribution (Fig. 16B). Only known from Campeche State in Mexico.

Nilothauma reissi (Soponis, 1987)

Figure 17C

Additional material. 1 male, slide-mounted: BRAZIL, Santa Catarina, São Francisco do Sul, Distrito do Saí, 26°13'40"S, 48°40'50"W, CEPA Vila da Glória, 11–15.xi.2019, #143, light trap, L.C. Pinho et al. leg. 1 male, slide-mounted: BRAZIL, São Paulo, Santa de Rosa Viterbo, bridge at Tio Zito, 27.ix. 2000, light trap, H.F. Mendes & T. Andersen leg. 1 male, slide-mounted: BRAZIL, Mato Grosso, Ribeirão Cascalheira, Estrada Fazenda Manaus, 1° af. Rio Bonito, 12°57.088'S, 51°52.480'W, 08.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg.

Distribution (Fig. 17C). The species was described from the Amazonas by Soponis (1987) and was later recorded from Minas Gerais and São Paulo States in northern and south-eastern Brazil by Mendes and Andersen (2009). The range is now extended to Mato Grosso and Santa Catarina States.

Nilothauma soka Andersen, Bello-González & Hagenlund, 2016

Figure 17C

Additional material. 2 males, slide-mounted: BRAZIL, Rondônia, Candeias do Jamari, Rio Preto, Ponte de Madeira, #01, 08°52'40"S, 63°38'02"W, 19–20.vii.2012, light trap, R. Boldrini & A.S. Fernandes leg. 2 males, slide-mounted: BRAZIL, Roraima, Boa Vista, Rio Cauamé, 02°52'06"N, 60°44'24"W, 9.iii.2009, light trap, L.M. Fusari

leg. 2 males, slide-mounted: BRAZIL, Amazonas, Barcelos, Rio Aracá, #9, 69 m a.s.l., 00°24'39"N, 63°23'12"W, 28.vii–06.viii.2009, light trap #3, N. Hamada et al. leg. 3 males, slide-mounted: BRAZIL, Amazonas, Barcelos, Rio Aracá, Foz do Igarapé Cuieiras, 00°19'15"N, 63°16'15"W, 35 m a.s.l., 30.vii–01.viii.2009, light trap #11, N. Hamada et al. leg.

Distribution (Fig. 17C). The species was originally described from the Amazonas State by Andersen et al. (2016); the range is now extended to the Rondônia and Roraima States in the Brazilian Amazon.

Nilothauma strebulosum (Adam & Sæther, 2000)

Figure 16D

Additional material. 1 male, slide-mounted: BRAZIL, Mato Grosso, Nova Xavantina, Fazenda Sr. Queté, Córrego Voadeira, 14°32.187'S, 52°30.902'W, 16.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg.

Distribution (Fig. 16D). The species was originally described from Costa Rica by Adam and Sæther (2000); the range is now extended to Mato Grosso State, central Brazil.

Nilothauma terena sp. nov.

<http://zoobank.org/140D5884-DBE1-403A-AD55-6AD631840573>

Figures 11A, B, 12A–H, 16A

Type material. Holotype male with larval and pupal exuvia, slide-mounted: BRAZIL, São Paulo, São Carlos, Campus UFSCar, Córrego do Fazzari, 21°59'S, 47°54'W, 11.ix.2008, L.C. Pinho & F.L. Silva leg. (UFSC).

Etymology. The specific epithet honours the Terena indigenous people from São Paulo State (Brazil). The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by having tergite IX with broadly rounded posterior margin without anal point, with anterolateral thorns, with dorsolateral lobes with few, strong setae and posteriolateral, narrowly subtriangular projection. The pupa can be recognised by having long, taeniate frontal setae and sternite I with extensive shagreen. The larva can be recognised by apparently having antenna with five segments only and by having mentum and inner teeth of mandible somewhat darker pigmented.

Description. Male imago (n = 1). Total length 2.67 mm. Wing length 1.21 mm. Total length/wing length 2.21. Wing length/length of profemur 2.11.

Colouration. Thorax and legs brown, abdomen light brown. Wing membrane without dark markings.

Antenna. AR = 0.21. Thirteenth flagellomere 120 µm long.




Figure 11. *Nilothauma terena* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with d tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Head. Temporal setae 5 in single row. Clypeus with 13 setae. Tentorium 62 μm long, maximum width 15 μm . Stipes not measurable. Palp segment lengths (in μm): 20, 23, 57, 92, 106. Third palpomere with 2 sensilla clavata subapically, longest about 25 μm . Fifth palpomere/third palpomere 1.87.

Thorax. Dorsocentrals 5 in single row, acrostichals 4, prealars 2. Scutellum with 2 setae.

Wing. VR = 1.46. Brachiolum with 1 seta, R with 6 setae, R_1 with 1 seta, R_{4+5} with 1 apical seta, remaining veins bare.

Legs. Spur of fore tibia 62 μm long including 32 μm long scale. Mid-leg missing; hind tibia with 2 spurs, 28 and 46 μm long. Combs of hind tibia 18 μm long. Width at apex of fore tibia 37 μm , of hind tibia 39 μm . Lengths and proportions of legs as in Table 10.

Hypopygium (Fig. 11A, B). Tergite IX with rounded posterior margin with altogether 6 marginal setae in two posteriolateral groups; with dorsolateral lobes with

Table 10. Lengths (in μm) and proportions of leg segments in *Nilothauma terena* sp. nov., adult male ($n = 1$).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	572	359	–	–	–	–
P ₂	–	–	–	–	–	–
P ₃	563	574	310	155	147	90
	ta ₅	LR	BV	SV	BR	
P ₁	–	–	–	–	–	
P ₂	–	–	–	–	–	
P ₃	49	0.57	3.22	3.58	7.1	

5 strong setae, longest setae 65 μm long; with posteriolateral, narrowly subtriangular projections, 14 μm long, 6 μm wide at base. Tergite band not apparent. Laterosternite IX without setae; with small anteriolateral thorn. Phallapodeme 33 μm long; transverse sternapodeme 11 μm long. Gonocoxite 83 μm long. Inferior volsella digitiform, curved, 55 μm long, 11 μm wide medially, with microtrichia and 12 short setae subapically. Superior volsella 29 μm long, 12 μm wide at base, 13 μm wide medially, covered with microtrichia and with marginal setae. Median volsella subtriangular with 2 apical setae on small tubercles, setae about 6 μm long. Gonostylus weakly curved, 97 μm long, 14 μm wide medially. HR = 0.85. HV = 2.75.

Female adult. Unknown.

Pupa ($n = 1$). Total length 3.59 mm. Exuviae pale brown.

Cephalothorax (Fig. 12A, B). Frontal apotome (Fig. 12A) with few wrinkles, frontal setae taeniate, 154 μm long. Thoracic horn not discernible; basal ring oval, 13 μm in diameter. Scutum with field of few weak tubercles. Anteprenotals 2; precorneals 2; dorsocentrals 4, Dc1 39 μm in front of Dc2, Dc2 96 μm in front of Dc3, Dc3 23 μm in front of Dc4.

Abdomen (Fig. 12C–E). Tergite I bare; tergites II–VI with transverse anterior band of somewhat stronger spinules, merging with median field of finer shagreen; anterior band of shagreen on tergite VI separated from posterior shagreen patch; tergite VII with anterior and posterior shagreen patches; tergite VIII with anterior shagreen patch connected with narrow posterior field of finer shagreen; tergite IX bare. Sternite I (Fig. 12D) with extensive shagreen; sternite II–VII bare; sternite VIII with narrow, longitudinal field of fine shagreen. Tergite II with 159 μm long row of 36 hooks, each hook about 8 μm long. Conjunctives III/IV and IV/V with spinules extending on to preceding segment. Pedes spurii B weakly developed on segment II. Anal comb 51 μm long, consisting of 3 spurs.

Abdominal setation. Lateral setae on segments I–VIII as: 0, 3, 3, 3, 4, 4, 4, 4; posterior lateral seta on tergite IV and all lateral setae on tergites V–VIII taeniate, remaining setae hair-like. All tergites with 1 pair of O setae.

Anal lobe. As long as broad, with 1 taeniate dorsal setae and complete fringe of 19 taeniae on each side, longest 170 μm . Male genital sac over-reaches anal lobe by 119 μm .

Fourth instar larva ($n = 1$). Head capsule 228 μm long. Postmentum 145 μm long.




Figure 12. *Nilothauma terena* sp. nov. pupa (A-E) and larva (F-H) A frontal apotome B thorax C abdomen, dorsal view D sternite E paratergite VIII F antenna G mandible H mentum and ventromental plates.

Head. Antenna (Fig. 12F) apparently with five segments only, length of antennal segments (in μm): 19, 10, 4, 11, 6. AR = 0.61. Basal antennal segment 10 μm wide; blade 32 μm long; accessory blade about 5 μm long. Premandible not measurable, teeth not discernible. Mandible (Fig. 12G) 69 μm long, seta subdentalis 19 μm long, inner teeth somewhat darker pigmented. Mentum (Fig. 12H) somewhat darker pigmented, 41 μm wide; middle part 10 μm wide with 2 minute inner teeth and pair of slightly larger lateral teeth; with 6 pairs of pointed, medially curved lateral teeth. Ventromental plates 98 μm wide, medially separated by 10 μm . Seta submenti 46 μm long.

Abdomen. Lost.

Distribution (Fig. 16A). Known from São Paulo State, south-eastern Brazil.

***Nilothauma txukuyana* sp. nov.**

<http://zoobank.org/8D8DBE02-E551-48F2-AFD8-DFF0B4263FCE>

Figures 13A, B, 16B

Type material. *Holotype* male, slide-mounted: BRAZIL, Pará, Rio Paru do Oeste, Malloca Apicó, 20.iv.1962, at light, E.J. Fittkau leg. (A 366-1, ZSM). *Paratypes*: 15 males, same data as holotype (ZSM, ZMBN, UFSC).

Etymology. The specific epithet honours the Txukuyana, indigenous people from Amazonas and Pará States in Brazil and from Suriname. The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by having tergite IX with broadly-rounded posterior margin without anal point, with dorso-lateral lobes with few, strong setae and posteriolateral, strongly setose, subtriangular projection.

Description. Male imago (n = 5–8). Total length 2.17–2.44, 2.25 mm. Wing length 1.00–1.09, 1.05 mm. Total length/wing length 2.02–2.33, 2.16. Wing length/length of profemur 2.16–2.26, 2.21.

Colouration. Head, thorax and legs brown; abdomen light brown. Wing membrane without dark markings.

Antenna. AR = 0.26–0.29, 0.28. Thirteenth flagellomere 132–156, 145 μm long.

Head. Temporal setae 6–10, 7 in single row. Clypeus with 13–16, 15 setae. Tentorium 69–83, 77 μm long, maximum width 14–19, 19 μm . Stipes not measurable. Palp segment lengths (in μm): 18–25, 21; 23–28, 26; 56–60, 57; 76–81, 79; 102–115, 107. Third palpomere with 2 sensilla clavata subapically, longest about 15 μm long. Fifth palpomere/third palpomere 1.72–1.98, 1.85.

Thorax. Dorsocentrals 4–7, 6 in single row, acrostichals apparently 4–6, 5 anterior, prealars 1–2, 2. Scutellum with 2 setae.

Wing. VR = 1.44–1.54, 1.50. Brachiolum with 1 seta, R with 7–9, 8 setae, R₁ with 5–8, 7 setae, R₄₊₅ with 9–13, 11 setae apically, remaining veins bare.

Legs. Spur of fore tibia 48–54, 52 μm long including 20–25, 22 μm long scale. Mid-tibia with 1 spur, 29–35, 32 μm long; hind tibia with 2 spurs, 22–28, 25 and




Figure 13. *Nilothauma txukuyana* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

33–39, 36 μm long. Combs of mid-tibia 17–18, 18 μm long, of hind tibia 19–22, 21 μm long. Width at apex of fore tibia 33–37, 35 μm , of mid-tibia 33–38, 36 μm , of hind tibia 39–41, 40 μm . Lengths and proportions of legs as in Table 11.

Hypopygium (Fig. 13A, B). Tergite IX with rounded posterior margin with 4–7, 5 marginal setae; with dorsolateral lobes with 3–4, 4 strong setae, longest setae 44–55, 50 μm long; with posteriolateral subtriangular, strongly setose projection, 35–41, 38 μm long, 22–25, 24 μm wide at base. Tergite band not apparent. Laterosternite IX with single setae. Phallapodeme 47–55, 49 μm long; transverse sternapodeme 17–19, 18 μm long. Gonocoxite 83–89, 87 μm long. Inferior volsella digitiform, curved, 62–72, 67 μm long, 11–14, 12 μm wide medially, with microtrichia and 13–17, 15 setae subapically. Superior volsella 25–29, 27 μm long, 8–11, 10 μm wide at base, 7–10, 9 μm wide medially, covered with microtrichia and with marginal setae. Median volsella small, broadly triangular, apparently without setae. Gonostylus nearly straight, 126–135, 130 μm long, 17–19, 18 μm wide medially. HR = 0.65–0.68, 0.66. HV = 1.78–1.87, 1.82.

Table 11. Length (in μm) and proportions of legs of *Nilothauma txukuyana* sp. nov., adult males (n = 5–7).

	Fe	Ti	ta ₁	ta ₂
P ₁	457–523, 482	319–359, 338	474–547, 515	194–221, 203
P ₂	449–474, 462	286–310, 301	147–179, 162	65–74, 69
P ₃	507–556, 529	458–482, 467	245–278, 263	114–139, 127
	ta ₃	ta ₄	ta ₅	LR
P ₁	147–163, 157	106–123, 114	57–65, 59	1.49–1.59, 1.52
P ₂	41–49, 46	25–33, 29	24–32, 26	0.50–0.58, 0.54
P ₃	114–147, 129	73–90, 78	41–49, 47	0.54–0.58, 0.56
	BV	SV	BR	
P ₁	2.47–2.54, 2.51	1.54–1.64, 1.59	2.43–3.46, 2.71	
P ₂	5.13–5.73, 5.45	4.36–5.06, 4.73	2.77–3.76, 3.38	
P ₃	3.20–3.57, 3.30	3.70–4.00, 3.79	4.33–5.00, 4.60	

Female imago and immatures. Unknown.

Distribution (Fig. 16B). Known from Pará State, Brazil.

***Nilothauma werekena* sp. nov.**

<http://zoobank.org/2B7C1444-B834-4FAD-BE69-E0417DD42812>

Figures 14A–C, 16D

Type material. *Holotype* male, slide-mounted: BRAZIL, Amazonas, Barcelos, Rio Aracá, Foz do Igarapé Cuieiras, #11, 00°19'15"N, 63°16'15"W, 35 m a.s.l., 30.vii–01.viii.2009, light trap, N. Hamada et al. leg. (UFSC). *Paratypes*: 4 males, slide-mounted, same data as holotype (INPA). 2 males, slide-mounted, same data as previous, except: #9, 00°24'39"N, 63°23'12"W, 69 m a.s.l., 28.vii–06.viii.2009, light trap #3, N. Hamada et al. leg. (MZSP).

Etymology. The specific epithet honours the Werekena indigenous people from the Rio Negro Basin in the Amazon. The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX with one low, but wide median dorsal protruberance with about 30 strong setae; anal point spatulate; superior volsella covered with microtrichia, fused to median volsella; laterosternite IX with thorn.

Description. Male imago (n = 5–7, unless otherwise stated). Total length 1.89–2.26, 2.11 mm. Wing length 0.98–1.11, 1.05 mm. Total length/wing length 1.90–2.15, 1.99. Wing length/length of profemur 2.21–2.53, 2.37.

Colouration. Head, thorax, legs and abdomen uniformly brown. Wing membrane without dark markings.

Antenna. AR = 0.32–0.39, 0.35. Thirteenth flagellomere 230–274, 260 μm long.

Head. Temporal setae 7–8, 7 in single row. Clypeus with 10–14, 12 setae. Tentorium 47–75, 65 μm long, maximum width 12–20, 17 μm . Stipes 80–117, 100 (4) μm long. Palp segment lengths (in μm): 20–32, 27; 22–27, 25; 65–85, 72; 95–125, 110; 87–132, 115. Third palpomere with 2–5, 4 sensilla clavata subapically, longest 12–15, 14 μm long. Fifth palpomere/third palpomere 1.21–1.75, 1.54.




Figure 14. *Nilothauma werekena* sp. nov. adult male **A** hypopygium, dorsal view **B** superior volsella, dorsal view **C** hypopygium with anal point and tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Thorax. Dorsocentrals 6–8, 7 in single row, acrostichals 10–14, 12, prealars 2. Scutellum with 2 setae.

Wing. VR = 1.21–1.32, 1.27. Brachiolum with 1 seta, R with 11–12, 11 setae, R_1 with 5–8, 7 setae, R_{4+5} with 11–17, 15 setae, remaining veins bare.

Legs. Spur of fore tibia 39–44, 42 μm long including 15–20, 17 μm long scale. Mid-tibia with 1 spur, 20–25, 22 μm long; hind tibia with 2 spurs, 20–25, 23 and 28–31, 30 μm long. Combs of mid-tibia 17–19, 18 μm long, of hind tibia 18–21, 19 μm long. Width at apex of fore tibia 29–39, 34 μm , of mid-tibia 34–39, 37 μm , of hind tibia 34–44, 39 μm . Lengths and proportions of legs as in Table 12.

Hypopygium (Fig. 14A–C). Tergite IX without dorsal lobe(s), with low, but wide median dorsal protruberance with 29–32, 31 strong, clustered median setae; posterior margin rounded to subrectangular, with 8–11, 9 weak setae to each side of base of anal point. Anal point spatulate, 37–47, 40 μm long, maximum width 5–10, 7 μm . Tergite bands not continuous. Laterosternite IX with 1–2, 1 seta; with anterolateral thorn. Phallapodeme 40–50, 45 μm long; transverse sternapodeme 15–25, 20 μm long. Gonocoxite 62–82, 72 μm long. Inferior volsella strongly curved, 22–30, 27 μm long,

Table 12. Lengths (in μm) and proportions of leg segments in *Nilothauma werekena* sp. nov., adult males (n = 5–7, unless otherwise stated).

	Fe	Ti	ta ₁	ta ₂
P ₁	374–501, 433	315–394, 345	443–522, 473	246–286, 266
P ₂	345–463, 424	296–345, 325	177–207, 197	89–99, 94
P ₃	443–532, 493	463–522, 502	266–305, 286	138–158, 148
	ta ₃	ta ₄	ta ₅	LR
P ₁	187–217, 207	118–148, 138 (4)	69–89, 79 (4)	1.36–1.47, 1.40
P ₂	69–79, 74	39–49, 44	30–39, 35	0.56–0.63, 0.60
P ₃	128–158, 148 (4)	89–99, 94 (4)	59–69, 55 (4)	0.57–0.61, 0.59
	BV	SV	BR	
P ₁	1.84–2.20, 1.95	1.55–1.66, 1.60	2.0–3.3, 2.5	
P ₂	3.50–4.25, 3.81	3.42–4.05, 3.71	2.3–4.7, 3.3	
P ₃	2.70–2.89, 2.81 (4)	3.39–3.45, 3.41	5.0–7.0, 5.9	

5–8, 6 μm wide medially, with microtrichia and 3 strong, simple setae apically. Superior volsella tongue-shaped to slightly pediform, 32–40, 37 μm long, 15–25, 20 μm wide at base, densely covered with microtrichia. Median volsella fused to superior volsella, consisting of 2–3, 2 small tubercles, each bearing single, long seta. Gonostylus 70–100, 90 μm long, basal half curved, distal half straight. HR = 0.63–0.96, 0.83. HV = 2.26–2.70, 2.34.

Female adult and immatures. Unknown.

Distribution (Fig. 16D). Known from Barcelos (Amazonas State), in the Brazilian Amazon.

Nilothauma yekwana sp. nov.

<http://zoobank.org/2FDCF2C9-C8ED-4675-93F2-65B893098864>

Figures 15A, B, 16A

Type material. *Holotype* male, slide-mounted: BRAZIL, Roraima, Boa Vista, BR-174, Igarapé Água Boa, 02°43'32"N, 60°48'43"W, 2014, N. Hamada leg (UFSC).

Etymology. The specific epithet honours the Ye'kwana, indigenous people from the Roraima State, Brazil. The name is to be regarded as a noun in apposition.

Diagnostic characters. The male can be distinguished from its congeners by the combination of: tergite IX with two setose dorsolateral lobes; anal point absent; posterior margin of tergite IX subrectangular; inferior volsella straight, tapering to apex; superior volsella curved, projecting posteriolaterally; median volsella broad, triangular, bearing 7 strong setae.

Description. Male imago (n = 1). Total length 1.77 mm. Wing length 0.91 mm. Total length/wing length 1.95. Wing length/length of profemur 2.19.

Colouration. Head, thorax, legs and abdomen uniformly light brown. Wing membrane without dark markings.

Antenna. AR = 0.22. Thirteenth flagellomere 137 μm long.

Head. Temporal setae 7 in single row. Clypeus with 15 setae. Tentorium 65 μm long, maximum width 12 μm . Stipes 80 μm long. Palp segment I–III lengths (in μm):




Figure 15. *Nilothauma yekwana* sp. nov. adult male **A** hypopygium, dorsal view **B** hypopygium with tergite IX removed, dorsal aspect to the left and ventral aspect to the right.

Table 13. Lengths (in μm) and proportions of leg segments in *Nilothauma yekwana* sp. nov., adult male ($n = 1$).

	Fe	Ti	ta ₁	ta ₂	ta ₃	ta ₄
P ₁	414	296	–	–	–	–
P ₂	394	256	177	59	39	30
P ₃	443	414	217	99	99	79
	ta ₅	LR	BV	SV	BR	
P ₁	–	–	–	–	–	–
P ₂	30	0.57	5.25	3.67	2.0	–
P ₃	49	0.52	3.60	3.95	4.4	–

22, 17, 50; segment IV and V lost. Third palpomere with 2 sensilla clavata subapically, longest 15 μm .

Thorax. Dorsocentrals 5 in single row, acrostichals 10, prealars 2. Scutellum with 2 setae.

Wing. VR = 1.48. Brachiolum with 1 seta, R with 6 setae, R₄₊₅ with 2 setae at apex, remaining veins bare.

Legs. Spur of fore tibia 44 μm long including 15 μm long scale. Mid-tibia with 1 spur, 20 μm long; hind tibia with 2 spurs, 20 and 25 μm long. Combs of mid-tibia 15 μm long, of hind tibia 18 μm long. Width at apex of fore tibia 39 μm , of mid-tibia 39 μm , of hind tibia 44 μm . Lengths and proportions of legs as in Table 13.




Figure 16. Distribution maps of Neotropical *Nilothauma* species **A** *N. hamadae* sp. nov., *N. jupau* sp. nov., *N. terena* sp. nov., *N. yekwana* sp. nov., *N. complicatum* Mendes & Andersen, 2009 **B** *N. leccii* sp. nov., *N. marianoii* sp. nov., *N. mateusi* sp. nov., *N. txukuyana* sp. nov., *N. maya* sp. nov. **C** *N. karitiana* sp. nov., *N. duena* Roback, 1960, *N. calori* Mendes & Andersen, 2009 **D** *N. werekena* sp. nov., *N. zitoi* Mendes & Andersen, 2009, *N. strebulosum* (Adam & Sæther, 2000).

Hypopygium (Fig. 15A, B). Tergite IX with 2 dorsolateral, densely setose lobes, setae about 15 μm long; with 2 strong setae anterolaterally, 2 medially and 4 close to posterior margin; posterior margin subquadrangular, anal point absent. Tergite bands lacking. Laterosternite IX with 1 seta. Phallapodeme 37 μm long; transverse sternapodeme 10 μm long. Gonocoxite 65 μm long. Inferior volsella straight, tapering to apex, 40 μm long, 7 μm wide medially, with microtrichia and 9 simple setae subapically. Superior volsella curved, projecting posteriolaterally, 22 μm long, 5 μm wide at base, covered with microtrichia and fringed at apex. Median volsella broad, triangular, 15 μm long, with 7 strong setae (one of them bifid), longest 20 μm . Gonostylus 75 μm long, straight. HR = 0.87. HV = 2.36.




Figure 17. Distribution maps of Neotropical *Nilothauma* species **A** *N. fittkai* (Soponis, 1987), *N. amazonense* Mendes & Andersen, 2009 **B** *N. aripuanense* Mendes & Andersen, 2009, *N. aleta* Roback, 1960 **C** *N. reissi* (Soponis, 1987), *N. soka* Andersen, Bello-González & Hagenlund, 2016 **D** *N. anamariae* Dantas & Hamada, 2017, *N. jaraguense* Mendes & Andersen, 2009.

Female adult and immatures. Unknown.

Distribution (Fig. 16A). Known from Roraima State, Brazilian Amazon.

***Nilothauma zitoi* Mendes & Andersen, 2009**

Figure 16D

Additional material. 1 male, slide-mounted: BRAZIL, Mato Grosso, Ribeirão Cascaheira, Fazenda Campina Verde, Rio Suiá Miçu, 12°48.591'S, 52°06.925'W, 10.x.2007, light trap, L.C. Pinho, S. Mateus, L. Torati & F.R. Silva leg.

Distribution (Fig. 16D). The species was originally described by Mendes and Andersen (2009), based on a single male from São Paulo State; the range is now extended to the Mato Grosso State.

Key to the males of *Nilothauma* Kieffer of the world

Modified from Qi et al. (2014), Niitsuma (2016) and Andersen et al. (2016), with the inclusion of sixteen species.

- 1 Tergite IX without setose dorsal lobe(s) or projection(s)..... 2
- Tergite IX with one to four setose dorsal lobes or projection(s) (e.g. Figs 16, 20) 19
- 2 Anal point present..... 3
- Anal point absent..... 15
- 3 Tergite IX with median cluster of about 30 strong setae..... 4
- Tergite IX with few, clustered setae, if numerous they are scattered (as in *N. aripuanense*) 5
- 4 Superior volsella slender, without microtrichia and with lateral spine. Brazil *Nilothauma jaquei* Dantas & Hamada, 2017
- Superior volsella pediform to lingulate, covered with microtrichia and without lateral spine (Fig. 14). Brazil..... *Nilothauma werekena* sp. nov.
- 5 Wing with conspicuous dark markings (Fig. 5); abdominal tergites II, III, and VI–VIII dark brown. Brazil..... *Nilothauma jupau* sp. nov.
- Wing unmarked, at most with faint colour (as in *Nilothauma aleta*, best seen in dark-field filter); abdominal tergites uniformly pale to brown 6
- 6 Gonostylus stout or swollen (Figs 1, 3, 6, 7) 7
- Gonostylus slender (Fig. 10) 10
- 7 Gonostylus very long, narrow basally and apically, swollen at mid-length (Fig. 6). Brazil *Nilothauma karitiana* sp. nov.
- Gonostylus stout, not distinctly swollen at mid-length (Figs 1, 3) 8
- 8 Acrostichals absent; anal point wide, covering most setae along posterior margin of tergite IX (Fig. 1). Peru, Brazil *Nilothauma aleta* Roback, 1960
- Acrostichals present; anal point comparatively narrow, nearly parallel-sided, with most setae placed lateral to base of anal point..... 9
- 9 Superior volsella tapering to apex; inferior volsella short, stout, with short, simple setae (Fig. 3). Peru, Brazil..... *Nilothauma duena* Roback, 1960
- Superior volsella wider at mid-length; inferior volsella long and slender, with long simple or apically split setae (Fig. 7). Brazil *Nilothauma leccii* sp. nov.
- 10 Superior volsella narrow, straight, curved or weakly sinuous, projecting posterior-medially, with one to six apical setae 11
- Superior volsella wider in distal half, projecting posterior-medially or posterior-laterally, with microtrichia only..... 13

- 11 Tergite IX with numerous scattered setae; anal point broadly lanceolate, about 20 μm wide. Brazil.....*Nilothauma aripuanense* Mendes & Andersen, 2009
- Tergite IX with one to four median setae and about 12 setae along posterior margin; anal point comparatively narrow 12
- 12 Tergite IX with single seta anterior to anal point; gonostylus with two to four sub-basal dorsal setae not arising from protuberances. Brazil.....
-*Nilothauma paucisetis* Dantas & Hamada, 2017
- Tergite IX with four aligned setae anterior to anal point; gonostylus with single sub-basal seta arising from distinct inner protuberance (Fig. 10). Neotropical Mexico..... *Nilothauma maya* sp. nov.
- 13 Anal point parallel-sided, about 4 μm wide; inferior volsella with about 18 slender, simple setae (Fig. 8). Brazil.....*Nilothauma marianoi* sp. nov.
- Anal point spatulate, 15–23 μm wide; inferior volsella with few (about 3) simple, slender setae or numerous (about 12) stout, split setae apically..... 14
- 14 Superior volsella boot-shaped, projecting posterior-laterally; inferior volsella narrow, with few simple, slender setae apically. Brazil.....
-*Nilothauma soka* Andersen, Bello-González & Hagenlund, 2016
- Superior volsella straight, projecting posterior-medially; inferior volsella wide, with numerous stout, split setae apically. Brazil.....
- *Nilothauma anamariae* Dantas & Hamada, 2017
- 15 Inferior volsella branched subapically. Brazil.....
-*Nilothauma complicatum* Mendes & Andersen, 2009
- Inferior volsella simple 16
- 16 Superior volsella pediform, without ventral transverse fold, with setae and microtrichia 17
- Superior volsella diamond-shaped, with ventral transverse fold, with microtrichia only 18
- 17 Wing vein R_1 with setae; gonostylus nearly parallel-sided in apical half. Brazil, Ecuador*Nilothauma fittkaui* (Soponis, 1987)
- Wing vein R_1 bare; gonostylus widest in apical one-third. Brazil.....
-*Nilothauma reissi* (Soponis, 1987)
- 18 Apex of superior volsella projecting caudad. Brazil.....
-*Nilothauma sooretamense* Mendes & Andersen, 2009
- Apex of superior volsella projecting mesad. Brazil.....
-*Nilothauma involucrum* Mendes & Andersen, 2009
- 19 Anal point lacking or rudimentary, completely covered by microtrichia 20
- Anal point present, with microtrichia at most in basal half..... 22
- 20 Tergite IX with four lobes or projections, one anterior pair with strong, long setae and one postero-lateral, triangular pair with weaker setae..... 21
- Tergite IX with one or two dorsal lobes..... 27

- 21 Tergite IX with postero-lateral pair of projections short, subequal in length to anterior pair (Fig. 11); laterosternite IX with thorn. Brazil*N. terena* **sp. nov.**
 – Tergite IX with postero-lateral pair of projections long, more than three times longer than anterior pair (Fig. 13); laterosternite IX without thorn. Brazil
 *N. txukuyana* **sp. nov.**
- 22 Dorsal projections of tergite IX differ in shape. Ghana.....
*Nilothauma insolitum* **Adam & Sæther, 1999**
 – Dorsal projections of tergite IX of the same shape.....**23**
- 23 Median volsella fused to superior volsella; superior volsella broadly pediform. Brazil *Nilothauma fazzariense* **Mendes & Andersen, 2009**
 – Median volsella distinct and separated from superior volsella; superior volsella digitate, curved, with or without lateral spine**24**
- 24 Dorso-lateral projections of tergite IX overreaching posterior margin of tergite. Brazil *Nilothauma roquei* **Mendes & Andersen, 2009**
 – Dorso-lateral projections of tergite IX not extended beyond posterior margin of tergite**25**
- 25 Superior volsella with lateral spine; laterosternite IX with thorn; posterior margin of tergite IX broadly rounded. Brazil.....
 *Nilothauma calori* **Mendes & Andersen, 2009**
 – Superior volsella without lateral spine; laterosternite IX without thorn; posterior margin of tergite IX subrectangular.....**26**
- 26 Median volsella consisting of single, small tubercle bearing one seta; inferior volsella curved, not tapering to apex. Brazil, Costa Rica.....
 *Nilothauma strebulosum* **(Adam & Sæther, 2000)**
 – Median volsella broad, triangular, bearing 7 strong setae; inferior volsella straight, tapering to apex (Fig. 15). Brazil*Nilothauma yekwana* **sp. nov.**
- 27 Tergite IX with single, median setose dorsal lobe.....**28**
 – Tergite IX with two setose dorsal lobes.....**37**
- 28 Superior volsella without microtrichia.....**29**
 – Superior volsella covered with microtrichia**32**
- 29 Dorsal projection weakly developed and undivided; situated posteriorly on tergite IX, close to base of anal point. China, Japan, Thailand.....
*Nilothauma japonicum* **Niitsuma, 1985**
 – Dorsal projection well developed, divided or undivided, situated anteriorly on tergite IX, at some distance from base of anal point**30**
- 30 Dorsal projection three-pronged at apex, without setae. Ghana.....
*Nilothauma fuscina* **Adam & Sæther, 1999**
 – Dorsal projection bell-shaped, with numerous setae**31**
- 31 Superior volsella with single apical seta; anal point narrow. Brazil
 *Nilothauma matogrossense* **Mendes & Andersen, 2009**
 – Superior volsella with one dorsolateral and one apical spine-like seta; anal point broad. Ghana *Nilothauma duminola* **Adam & Sæther, 1999**

- 32 Anal point very broad, about half as wide as tergite IX and lanceolate (Fig. 4).
Brazil *Nilothauma hamadae* sp. nov.
- Anal point narrow, spatulate to tapering, but not lanceolate..... 33
- 33 Anal point tapering or only slightly widened medially..... 34
- Anal point distinctly spatulate..... 35
- 34 Wing with distinct dark areas at RM, FCu, along apical half of An and in cells r_{4+5}
and m_{1+2} , Cuba
..... *Nilothauma granma* Andersen, Bello-González & Hagenlund, 2016
- Wing without dark areas. Canada, USA..... *Nilothauma babyi* (Rempel, 1937)
- 35 Dorsal projection large, covering most of tergite IX. Australia.....
..... *Nilothauma adunatum* Adam & Sæther, 1999
- Dorsal projection of tergite IX small, with setae at apex only 36
- 36 Wing length > 2.4 mm; AR = 0.28; median volsella consisting of two tubercles,
each with single, strong apical seta. Canada.....
..... *Nilothauma verrucum* Adam & Sæther, 1999
- Wing length < 1.3 mm; AR = 0.13; median volsella consisting of three tubercles,
each with strong, apical seta. Venezuela.....
..... *Nilothauma canaima* Andersen, Bello-González & Hagenlund, 2016
- 37 Dorsal projections of tergite IX of the same shape..... 38
- Dorsal projections of tergite IX of different shapes 42
- 38 Anal point short, digitiform, with microtrichia in basal half; gonostylus distinctly
widened in apical one-third. Brazil.... *Nilothauma zitoi* Mendes & Andersen, 2009
- Anal point well developed, lanceolate or parallel-sided; gonostylus nearly parallel-
sided in apical half 39
- 39 Superior volsella short, subtriangular, with two apical tubercles bearing setae; in-
ferior volsella with long, strong simple setae (Fig. 9). Brazil.....
..... *Nilothauma mateusi* sp. nov.
- Superior volsella long, pediform to tongue-shaped; inferior volsella with slender,
simple apically split setae..... 40
- 40 Anal point parallel-sided; laterosternite IX with thorn. Chile
..... *Nilothauma spiesi* Mendes & Andersen, 2009
- Anal point lanceolate; laterosternite IX without thorn..... 41
- 41 Inferior volsella and gonostylus with apically split setae; median volsella curved,
tapering, with microtrichia and setae. Brazil.....
..... *Nilothauma jaraguaense* Mendes & Andersen, 2009
- Inferior volsella and gonostylus with simple setae only; median volsella short,
parallel-sided, with two apical setae, without microtrichia. Brazil.....
..... *Nilothauma amazonense* Mendes & Andersen, 2009
- 42 Wing with dark areas or bands; anterior projection on tergite IX long and deeply
divided..... 43
- Wing without dark markings; anterior projection on tergite IX variably devel-
oped..... 50

- 43 Anterior projection on tergite IX with setae not restricted to apex. South Africa .
..... *Nilothauma harrisoni* Adam & Sæther, 1999
- Anterior projection on tergite IX with apical setae only.....44
- 44 Posterior projection on tergite IX deeply divided into antero-dorsal and postero-ventral parts45
- Posterior projection on tergite IX not as above47
- 45 Anterior projection on tergite IX with simple, separated, apical setae; anterior part of posterior projection apically pointed; wing with postero-median spot extending on both sides of Cu_1 . Afrotropical
..... *Nilothauma pictipenne* Kieffer, 1921
- Anterior projection on tergite IX with setae forming fan-like structure; anterior part of posterior projection with blunt apex; wing with postero-median spot exclusively proximal of Cu_1 46
- 46 Wing with four dark areas; setae on anterior projection on tergite IX branched apically. Ghana.....*Nilothauma flabellatum* Adam & Sæther, 1999
- Wing with three dark areas; apical setae on anterior projection on tergite IX lamellate, not branched. Ghana..... *Nilothauma kakumense* Adam & Sæther, 1999
- 47 Posterior projection on tergite IX either with a disto-dorsal lobe or subapical constriction48
- Posterior projection on tergite IX without distal lobe or constriction49
- 48 Posterior projection on tergite IX with long antero-lateral arms and disto-dorsal lobe. Tanzania *Nilothauma anderseni* Adam & Sæther, 1999
- Posterior projection on tergite IX without antero-lateral arms, with apical constriction and five apical setae. Zimbabwe.....
..... *Nilothauma latocaudatum* Adam & Sæther, 1999
- 49 Posterior projection on tergite IX without microtrichia, except on long, antero-lateral arms; superior volsella without antero-median extension; median volsella present. China, Japan *Nilothauma nojirimaculatum* Sasa, 1991
- Posterior projection on tergite IX without antero-lateral arms, covered with microtrichia; superior volsella with antero-median extension; median volsella apparently absent. Japan *Nilothauma hibaraqartum* Sasa, 1993
- 50 Anterior projection on tergite IX with apically plumose setae.....51
- Apical setae on anterior projection on tergite IX not plumose58
- 51 Anal point with microtrichia along median ridge and apical margin. Oriental China.....*Nilothauma pandum* Qi, Lin, Wang & Shao, 2014
- Anal point without microtrichia.....52
- 52 Superior volsella pad-like, with extensive microtrichia. Palearctic Japan.....
..... *Nilothauma hibaratertium* Sasa, 1993
- Superior volsella slender, microtrichia absent or limited within small area when present53
- 53 Superior volsella without lateral spur. Thailand
..... *Nilothauma mergae* Adam & Sæther, 1999
- Superior volsella with lateral spur54

- 54 Superior volsella without setal brush or fringe, with only one to three apical setae **55**
- Superior volsella with apical setal brush or fringe **56**
- 55 Anal point with microtrichia along median ridge; superior volsella relatively long when compared to median volsella (length ratio, $Svo/Mvo > 4.0$); inferior volsella with simple setae only. Oriental China.....
..... ***Nilothauma aristatum* Qi, Tang & Wang, 2016**
- Anal point bare; length ratio Svo/Mvo about 2.0; inferior volsella with apically split setae. Oriental China.....***Nilothauma acre* Adam & Sæther, 1999**
- 56 Superior volsella four to five times as long as median volsella. Palaearctic Japan ..
.....***Nilothauma niidaensis* Niitsuma, 2016**
- Superior volsella two to three times as long as median volsella.....**57**
- 57 Anterior projection on tergite IX two to four times as long as broad. Canada, USA.....***Nilothauma bicorne* (Townes, 1945)**
- Anterior projection on tergite IX broader than long. USA.....
..... ***Nilothauma mirabile* (Townes, 1945)**
- 58 Anal point trifid; anterior projection on tergite IX very long, tapering to parallel-sided apex, with apical setae only; posterior projection on tergite IX distally very slender, with five apical setae. D. R. Congo, Ghana
..... ***Nilothauma burmeisteri* Adam & Sæther, 1999**
- Anal point simple; anterior projection on tergite IX wart-like, with setae not only at apex; posterior projection on tergite IX triangular or apically rounded **59**
- 59 Posterior projection on tergite IX apically rounded; superior volsella with two to four lobes..... **60**
- Posterior projection on tergite IX triangular; superior volsella without lobes ... **61**
- 60 Anterior projection on tergite IX partially divided apically; superior volsella with two lobes. Oriental China..... ***Nilothauma bilobatum* Qi, Tang & Wang, 2016**
- Anterior projection on tergite IX undivided apically; superior volsella with four lobes. Oriental China.... ***Nilothauma quatuorlobum* Yan, Tang & Wang, 2005**
- 61 Anal point parallel-sided; anterior projection on tergite IX with setae thickened at apices. Ghana..... ***Nilothauma ankasense* Adam & Sæther, 1999**
- Anal point spatulate; anterior projection on tergite IX with setae not thickened at apices **62**
- 62 Superior volsella tapering, widest near base. Europe
.....***Nilothauma brayi* (Goetghebuer, 1921)**
- Superior volsella widest about one-third from apex. Australia.....
.....***Nilothauma infissum* Adam & Sæther, 1999**

Acknowledgements

We are indebted to Dr. Neusa Hamada and all her team at INPA for providing rich material from the Amazon and to Dr. Martin Spies for the loan of material from the

Zoologische Staatssammlung München, Germany. Thanks are also due to Dr. Sidnei Mateus, Dr. Marcia R. Spies, Dr. Adolfo R. Calor and the staff at the Laboratory of Systematics of Diptera (UFSC) for all help and for providing us with freshly collected material. Fieldwork in Santa Catarina was funded by the Santa Catarina Research Foundation (FAPESC) – project “Insetos aquáticos do Parque Estadual da Serra Furada, SC” (FAPESC 11323/2012–9) and by the City Hall of São Francisco do Sul – project “Nascentes do Sai”.

References

- Adam JI, Sæther OA (1999) Revision of the genus *Nilothauma* Kieffer, 1921 (Diptera: Chironomidae). *Entomologica Scandinavica*, Supplement 56: 1–107. <https://doi.org/10.1051/limn/2016013>
- Adam JI, Sæther OA (2000) *Paranilothauma strebulosa* sp. nov. from Costa Rica (Diptera: Chironomidae). In: Hoffrichter O (Ed.) Late 20th Century Research on Chironomidae: An Anthology from the 13th International Symposium on Chironomidae. Shaker Verlag, Aachen: 19–23.
- Andersen T, González OCB, Hagenlund LK (2016) Three new Neotropical species of *Nilothauma* Kieffer, 1921 (Diptera: Chironomidae). *Annales de Limnologie – International Journal of Limnology* 52: 253–261. <https://doi.org/10.1051/limn/2016013>
- Dantas GPS, Hamada N (2017) Three new species of *Nilothauma* Kieffer (Diptera: Chironomidae) from Brazil. *Zootaxa* 4282: 350–360. <https://doi.org/10.11646/zootaxa.4282.2.8>
- Epler JH (1988) Biosystematics of the genus *Dicrotendipes* Kieffer, 1913 (Diptera: Chironomidae: Chironominae) of the world. *Memoirs of the American Entomological Society* 36: 1–214.
- Epler JH, Ekrem T, Cranston PS (2013) 10. The larvae of Chironominae (Diptera: Chironomidae) of the Holarctic Region – Keys and diagnoses. In: Andersen T, Cranston PS, Epler JH (Eds) The larvae of Chironomidae (Diptera) of the Holarctic Region – Keys and diagnoses. *Insect Systematics & Evolution*, Supplement 66: 387–556.
- Kieffer JJ (1921) Synopse de la tribu des Chironomariae (Diptères). *Annales de la Société scientifique de Bruxelles* 40(3–4): 269–277.
- Mendes HF, Andersen T (2009) Neotropical *Nilothauma* Kieffer, 1921, with the description of thirteen new species (Diptera: Chironomidae). *Zootaxa* 2063: 1–45. <https://doi.org/10.11646/zootaxa.2063.1.1>
- Niitsuma H (2016) Two species of *Nilothauma* Kieffer (Diptera, Chironomidae) from Japan, with description of a new species. *Zootaxa* 4079: 573–581. <https://doi.org/10.11646/zootaxa.4079.5.4>
- Pinho LC, Mendes HF, Andersen T (2009) New species and records of *Beardius* Reiss & Sublette from Brazil (Diptera, Chironomidae). *Spixiana* 32: 255–264.
- Pinho LC, Mendes HF, Andersen T (2013) Revision of *Beardius* Reiss et Sublette, 1985 (Diptera: Chironomidae), with the description of twenty new species. *Zootaxa* 3742: 1–78. <https://doi.org/10.11646/zootaxa.3742.1.1>

- Qi X, Lin X, Wang X, Shao Q (2014) A new species of *Nilothauma* Kieffer from China, with a key to known species of the genus (Diptera: Chironomidae). *Zootaxa* 3869: 573–578. <https://doi.org/10.11646/zootaxa.3869.5.7>
- Qi X, Tang H, Wang X (2016) Notes on *Nilothauma* Kieffer from Oriental China, with descriptions of three new species (Diptera, Chironomidae). *ZooKeys* 574: 143–159. <https://doi.org/10.3897/zookeys.574.6129>
- Roback SS (1960) Results of the Catherwood Foundation Peruvian Amazon Expedition. New species of Tendipedidae (Diptera). *Transactions of the American Entomological Society* 86: 87–107.
- Sæther OA (1980) Glossary of chironomid morphology terminology (Diptera: Chironomidae). *Entomologica Scandinavica, Supplement* 14: 1–51.
- Soponis AR (1987) *Paranilothauma* and *Neelamia*, new genera of Chironomini (Diptera: Chironomidae) from Brazil. *Studies on Neotropical Fauna and Environment* 22: 11–24. <https://doi.org/10.1080/01650528709360715>
- Tokeshi M (1995) Life cycles and population dynamics. In: Armitage PD, Cranston PS, Pinder LCV (Eds) *The Chironomidae: Biology and ecology of non-biting midges*. Chapman & Hall, London, 225–268. https://doi.org/10.1007/978-94-011-0715-0_10